arXiv:1507.03191v3 [math.PR] 20 Dec 2016

ON GENERALIZED KESTEN–MCKAY DISTRIBUTIONS

PAWEŁ J. SZABŁOWSKI

Abstract. We examine properties of distributions with the density of the form: $\frac{2A_n\sqrt{c^2-x^2}}{\pi c\prod_{j=0}^{n}(c(1+a_j^2)-2a_jx)}$, where $c,a_1,\ldots,a_n$ are some parameters and A_n suitable constant. We find general forms of A_n, of $k-$th moment and of $k-$th polynomial orthogonal with respect to this measure. We indicate their connections with polynomials forming the so called Askey–Wilson scheme. On the way we prove several identities concerning rational symmetric functions. Finally we consider the case of parameters $a_1,\ldots,a_n$ forming conjugate pairs and give multivariate interpretation of obtained distributions at least for the cases $n=2,4,6$.

1. Introduction

The purpose of this note is to analyze the properties of the following family of distributions having densities of the form:

$$f_{KMKn}(x|c,a_1,\ldots,a_n)=\frac{2A_nc^{n-2}\sqrt{c^2-x^2}}{\pi\prod_{j=1}^{n}(c(1+a_j^2)-2a_jx)}, \tag{1.1}$$

defined for $n\geq 0$, $|x|\leq c$, with $c>0$, $|a_j|<1$, $j=1,\ldots,n$. Here A_n is a normalizing constant being the function of parameters $a_1,\ldots,a_n$. We will call this family generalized Kesten–McKay distributions. The name is justified by the fact that the distribution with the following density

$$f_{KMK2}(x|4/a^2,a,-a)=\frac{v\sqrt{4(v-1)-x^2}}{2\pi(v^2-x^2)}, \tag{1.2}$$

where $v=1+1/a^2$ and $|a|<1$ has been defined, described and what is more, derived in [5] or [9]. Then the name Kesten-McKay distribution has been attributed to this distribution in the literature that appeared after 1981.

Thus it is justified to call the distribution defined by (1.1) a generalized Kesten-McKay (GKM) distribution.

Note also that for $n=0$ $f_{KMK0}(x|c)$ becomes Wigner or semicircle distribution with parameter c.

It should be underlined that for $n=0,1,2$ distributions of this kind appear not only in the context of random matrices, random graphs which is typical application of Kesten–Mckay distributions (see e.g.[7], [8], [10]) but also in the context of the

---

Date: June 2015.

2010 *Mathematics Subject Classification.* Primary 60E05,05E05; Secondary 62H05,60J10.

Key words and phrases. Kesten–McKay, Bernstein-Szegö distributions, Chebyshev polynomials, orthogonal polynomials, Askey–Wilson polynomials, moments, symmetric rational functions, multivariate distributions, .

The author is very grateful to the unknown referee for his more or less detailed remarks and questions that enabled improvement of the paper.

so-called free probability a part of non-commutative probability theory recently rapidly developing. One of the first papers where semicircle and related distribution appeared in the non-commutative probability context is [1].

For $n<5$ distribution f_{KMKn} can be identified as the special case of the Askey–Wilson chain of distribution that make 5 families of polynomials of the so-called Askey–Wilson scheme of polynomials, orthogonal. For the reference see e.g. [13] For $n\geq 5$ the distributions f_{KMKn} were not yet described in detail.

It has to be noted that in 2009 there appeared paper [3]. Although the aim of it was to analyze two dimensional measures on the plane of the form

$$\frac{\sqrt{(1-x^{2})(1-y^{2})}}{g(x,y)},$$

where g was a polynomial in x and y it also contains some results concerning $1-$dimensional case. The considered there $1-$ dimensional distribution is very much alike the distribution we consider in this paper. The authors of the paper call it Bernstein-Szegö distribution. The method they use to analyze these distributions allow them to consider only the case of even degree n of polynomial g. The results of the paper are general and hence rather imprecise. They were obtained by quite complicated integration on the complex plane. Our results are precise since we assume the precise form of the polynomial in the denominator. Namely we assume the knowledge of the roots of this polynomial. Besides as mentioned above the order of this polynomial can be also odd. Moreover due to our assumptions we are able to give precise form of the polynomials orthogonal with respect to this measure.

Out methods are simple and heavily exploit the properties of Chebyshev polynomials of the second kind allowing to obtain on the way some interesting identities like e.g. the ones given in Remark 3 or Corollary 2.

It should be mentioned that by considering polynomials of even degree, having pairwise conjugate roots we are not only able to cover the case of two dimensional measures on the plane of the form mentioned above but also generalize the results to 3 or more dimensions defying new distributions and finding they marginals. Compare Section 2 and the remarks at the end of this section.

Hence our results give substance and generalize the results of [3].

We will present unified approach and recall and collect information on this family that is scattered though literature.

Let us observe first that if $X\sim f_{KMKn}(x|c,a_{1},\ldots,a_{n})$ and $Y\sim f_{KMKn}(x|1,a_{1},\ldots,a_{n})$ then $X\sim cY.$ Hence we will consider further only distributions with the density $f_{KMKn}(x|1,a_{1},\ldots,a_{n})$ which we will denote for brevity $f_{Kn}(x|a_{1},\ldots,a_{n})$.

To start analysis let us recall that

$$1/(1+a_{j}^{2}-2a_{j}x)=\sum_{k=0}^{\infty}a_{j}^{k}U_{k}(x), \tag{1.3}$$

where U_{k} denotes $k-$th Chebyshev polynomial of the second kind and that

$$\int_{-1}^{1}\frac{2}{\pi}U_{k}(x)U_{j}(x)\sqrt{1-x^{2}}dx=\left\{\begin{array}{lll}0 & if & j\neq k\\ 1 & if & j=k\end{array}\right. .$$

Thus the form of the density (1.1) fits the scheme of distributions and orthogonal polynomials that was considered in [13] and hence we can use ideas and results presented there. First of all let us observe that the densities considered in this paper are the special cases of the distributions known from the so-called Askey–Wilson scheme of distributions and orthogonal polynomials obtained from the general ones by setting $q=0$. q is a special parameter called base in the theory of Askey–Wilson polynomials. More precisely for $n=0$ we deal with Wigner distribution, for n $=1$ we deal with the so-called continuous big $q-$Hermite polynomials and the distribution that makes them orthogonal. When we set $q=0$ and $n=1$ we deal with distribution $f_{K1}(x|a_1)$. For $n=2$ we deal with the so-called Al-Salam–Chihara polynomials and the distribution that make these polynomials orthogonal. Setting now $q=0$ leads us to distribution $f_{K2}(x|a_1,a_2)$. Further for $n=3$ we deal with the so-called dual Hahn polynomials and the distribution that make these polynomials orthogonal. Finally for $n=4$ we deal with the so-called Askey–Wilson polynomials and the distribution that make them orthogonal. Hence in particular we know the families of orthogonal polynomials that our distributions for $n=0,\ldots,4$ make orthogonal. For the precise definitions and further references see e.g. [6] or [13].

In the sequel we will use exchangeably $\mathbf{a}_n$ and $(a_1,\ldots,a_n)$ depending on the required brevity and clarity.

Let us denote $\prod_{j=1,j\neq i}^{n}$ as $\prod_{j\neq i}^{n}$.

We have the following general observation.

Theorem 1. *If* $a_i\neq a_j$, $i\neq j$, $i,j=1,\ldots,n$, *then*
i) constant A_n *in (1.1) is given by:*

$$A_n=A_n(\mathbf{a}_n)=1/\sum_{i=1}^{n}\frac{a_i^{n-1}}{\prod_{j\neq i}^{n}(a_i-a_j)(1-a_ia_j)}. \tag{1.4}$$

ii) Let us define constants $B_{n,k}$ *for* $k\geq 0$ *by:*

$$B_{n,k}(\mathbf{a}_n)=\int_{-1}^{1}U_k(x)f_{Kn}(x|\mathbf{a}_n)dx,$$

Then

$$B_{n,k}=A_n\sum_{i=1}^{n}\frac{a_i^{n+k-1}}{\prod_{j\neq i}^{n}(a_i-a_j)(1-a_ia_j)}, \tag{1.5}$$

for $k=0,\ldots.$

Proof. We will use the fact that

$$1/\prod_{i-1}^{n}(x-b_i)=\sum_{i=1}^{n}\frac{1}{\prod_{j=1,j\neq i}^{n}(b_i-b_j)}\frac{1}{(x-b_i)},$$

and the fact that $((1+a^2)/a-(1+b^2)/b)=(b-a)(1-ab)/(ab)$. Hence we have

$$\begin{aligned}
f_{Kn}(x|a_1,\ldots,a_n) &= \frac{2A_n(-1)^n}{2^n\pi\prod_{i=1}^n a_i}\sqrt{1-x^2}/\prod_{i=1}^n(x-(1+a_i^2)/(2a_i)) \\
&= \frac{2A_n(-1)^n}{2^n\pi\prod_{i=1}^n a_i}\sqrt{1-x^2}\sum_{i=1}^n\frac{a_i^{n-1}\prod_{j\neq i}^n a_j}{\prod_{j\neq i}^n(a_j-a_i)(1-a_ia_j)}\frac{1}{(x-(1+a_i^2)/(2a_i))} \\
&= \frac{2A_n(-1)^n}{\pi}\sqrt{1-x^2}\sum_{i=1}^n\frac{a_i^{n-1}}{\prod_{j\neq i}^n(a_i-a_j)(1-a_ia_j)}\frac{1}{((1+a_i^2)-2a_ix)}.
\end{aligned}$$

Now following (1.3) and orthogonality of polynomials $U's$ with respect to Wigner measure we have:

$$\int_{-1}^{1}\frac{2\sqrt{1-x^2}}{\pi((1+a_i^2)-2a_ix)}U_k(x)dx=a_i^k.$$

□

Remark 1. *Since coefficients $B_{n,k}$ are coefficients in the orthogonal expansion in $L_2([-1,1],w)$, where w denotes measure with the density $\frac{2}{\pi}\sqrt{1-x^2}$, in basis $\{U_k\}_{k\geq0}$, we get the following expansion for free:*

$$f_{Kn}(x|\mathbf{a}_n)=\frac{2}{\pi}\sqrt{1-x^2}\sum_{k=0}^{\infty}B_{n,k}U_k(x). \tag{1.6}$$

For more examples of such expansions see [11].

In the sequel we will need the following quantities:

$$\begin{aligned}
S_k^{(n)}(\mathbf{a}_n) &= \sum_{0\leq j_1<j_2<\ldots,<j_k\leq n}\prod_{m=1}^k a_{j_m} &(1.7)\\
\Delta_m^{(n)}(\mathbf{a}_n) &= \sum_{\substack{0\leq j_1,\ldots j_{n-1}\leq m\\ j_1+\ldots+j_{n-1}\leq m}} a_1^{j_1}a_2^{j_2}\ldots a_{n-1}^{j_{n-1}}a_n^{m-(j_1+\ldots+j_{n-1})}, &(1.8)
\end{aligned}$$

for $1\leq k\leq n\leq m$. Whenever it will be obvious we will drop other than k arguments of the functions S_k and Δ_k. Notice that $S_k^{(n)}$ is the $k-$th elementary symmetric function of the variables $a_1,\ldots,a_n$. We set $S_0^{(n)}(\mathbf{a}_n)=1$.

Remark 2. *We have*

$$\begin{aligned}
A_1 &= 1, A_2=1-a_1a_2,\ A_3=\prod_{j=1}^3\prod_{k=j+1}^3(1-a_ja_k)=(1-a_1a_2)(1-a_1a_3)(1-a_2a_3),\\
A_4 &= \prod_{j=1}^4\prod_{k=j+1}^4(1-a_ja_k)/(1-S_4)=\\
&\frac{(1-a_1a_2)(1-a_1a_3)(1-a_2a_3)(1-a_1a_4)(1-a_2a_3)(1-a_2a_4)(1-a_3a_4)}{1-a_1a_2a_3a_4},\\
A_5 &= \prod_{j=1}^5\prod_{k=j+1}^5(1-a_ja_k)/(1-S_4+S_1S_5-S_5^2),\\
A_6 &= \frac{\prod_{j=1}^6\prod_{k=j+1}^6(1-a_ja_k)}{(1-S_4+S_1S_5-S_5^2-S_6-S_1^2S_6+S_2S_6+S_4S_6+S_1S_5S_6-S_6^2-S_2S_6^2+S_6^3)}.
\end{aligned}$$

Notice that for $n = 1, \ldots, 4$ the A_n agrees with respective constants given in [13] *in formulae (2.4), (2.6) and unnamed formulae on top of 9-th and 10-th pages for $q = 0$, when presenting respectively densities that make big q-Hermite, Al-Salam–Chihara, dual Hahn and Askey-Wilson polynomials orthogonal.*

Using this denotations we have: $B_{2,k} = \Delta_k^{(2)}$, $B_{3,k} = \Delta_k^{(3)} - S_3\Delta_{k-1}^{(3)}$. Compare here with formulae (2.7) and (2.10) in [13] *for $q = 0$.*

Remark 3. *Let us notice also that*

$$\begin{aligned} 1/A_n(\mathbf{a}_n) &= \int_{-1}^{1} \frac{2}{\pi}\sqrt{1-x^2}\prod_{i=1}^{n}\sum_{k_i=1}^{\infty} a_i^{k_i} U_{k_i}(x)dx = \\ &\sum_{k_1\geq 0,\ldots,k_n\geq 0}(\prod_{j=1}^{n} a_i^{k_i})V_{k_1,\ldots,k_n}, \end{aligned}$$

where $V_{k_1,\ldots,k_n} = \int_{-1}^{1}\frac{2}{\pi}\sqrt{1-x^2}\prod_{i=1}^{n}U_{k_i}(x)dx$. That is we get generating function of the numbers $V_{k_1,\ldots,k_n}$ for free. This observation gives also interesting interpretation of constants A_n.

Corollary 1. *i)*

$$\int_{-1}^{1} x^k f_{Kn}(x|\mathbf{a}_n)dx = \frac{1}{(k+1)2^k}\sum_{j=0}^{\lfloor k/2\rfloor}(k-2j+1)\binom{k+1}{j}B_{n,k-2j}(\mathbf{a}_n),$$

ii)

$$\int_{-1}^{1} U_k(x)U_m(x)f_{Kn}(x|\mathbf{a}_n)dx = \sum_{j=0}^{\min(m,k)} B_{n,|m-k|+2j}(\mathbf{a}_n).$$

Proof. i) we used well known identity: $2^k x^k = \sum_{j=0}^{\lfloor k/2\rfloor}(\binom{k}{j} - \binom{k}{j-1})U_{k-2j}(x)$ (see e.g. [16] Proposition 1 with $q = 0$ and the fact that $h_n(x|0) = U_n(x)$) and the fact that $\binom{k}{j} - \binom{k}{j-1} = (k-2j+1)\binom{k+1}{j}/(k+1)$. ii) we used identity $U_k(x)U_m(x) = \sum_{j=0}^{\min(k,m)} U_{|k-m|+2j}(x)$ that can be easily derived from $2T_n(x)T_m(x) = T_{n+m}(x) + T_{|n-m|}(x)$, where T_n denotes Chebyshev polynomial of the first kind and the formulae relating Chebyshev polynomials of the first and second kind. See also formula (2.13) of [15] with $q = 0$. □

Proposition 1. *For any function $g : \mathbb{R} \longrightarrow \mathbb{R}$ let us denote $\mathbf{g}(\mathbf{a}_n) = (g(a_1), \ldots, g(a_n))$. Also let $\mathbf{b}_n^{(i)} = (b_1, \ldots, b_{i-1}, b_{i+1}, \ldots, b_n)$. Let $g(x) = (1+x^2)/(2x)$. Then if $a_i \neq a_j$, $i \neq j$, $i, j = 1, \ldots, n$, we have for every $k = 1, \ldots, n-1$:*

$$\sum_{i=1}^{n}\frac{a_i^{n-2}S_k(\mathbf{g}(\mathbf{a}_n^{(i)}))}{\prod_{j\neq i}^{n}(a_j - a_i)(1 - a_i a_j)} = 0. \tag{1.9}$$

Proof. We start from the fact that $1/\prod_{i-1}^{n}(x - b_i) = \sum_{i=1}^{n} D_{n,i}\frac{1}{(x-b_i)}$ where $D_{n,i} = 1/\prod_{j=1,j\neq i}^{n}(b_i - b_j)$. Thus from the properties of simple fractions decompositions we have the following identity

$$1/\prod_{i-1}^{n}(x - b_i) = (\sum_{i=1}^{n} D_{n,i}\prod_{j\neq i}(x - b_j))/\prod_{i=1}^{n}(x - b_i).$$

Hence all coefficients by nonzero powers of x must be zero. In particular for $\forall k = 0\ldots, n-1$ we must have $\sum_{i=1}^{n} D_{n,i} S_k(\mathbf{b}_n^{(i)}) = 0$. Now it remains to substitute $b_i = g(a_i)$ and use the fact that $(b_i - b_j) = (a_j - a_i)(1 - a_i a_j)/(2a_i a_j)$ so that

$$D_{n,i} = 2^{n-1} a_i^{n-2} (\prod_{j=1}^{n} a_j) / \prod_{j\neq i}^{n} ((1 - a_i a_j)(a_j - a_i))$$

since $a_i^{n-1} \prod_{j\neq i}^{n} a_j = a_i^{n-2} \prod_{j=1}^{n} a_j$. □

As a corollary we get in particular the following identities:

Corollary 2. $\forall n \geq 2$: *i)*

$$\sum_{i=1}^{n} \frac{a_i^{n-2}}{\prod_{j\neq i}^{n} (a_i - a_j)(1 - a_i a_j)} = 0, \tag{1.10}$$

ii)

$$\sum_{i=1}^{n} \frac{a_i^{n-2}(1 + a_i^2) \sum_{j\neq i} \prod_{k\neq i} a_k}{\prod_{j\neq i}^{n} (a_i - a_j)(1 - a_i a_j)} = 0,$$

iii) $\forall m \geq 1, n \geq 2$:

$$\sum_{j=0}^{n} (-1)^j S_j^{(n)}(\mathbf{a}_n) B_{n,m-j}(\mathbf{a}_n) = 0. \tag{1.11}$$

Proof. To prove i) we take $k = 0$ in (1.9). To prove ii) we take $k = 1$ in (1.9) and notice that

$$\begin{aligned} a_i^{n-1} (\prod_{j\neq i}^{n} a_j) S_1(\mathbf{g}(\mathbf{a}_n^{(i)}) &= a_i^{n-1} \sum_{j\neq i}^{n} (1 + a_j^2) \prod_{k\neq i,j}^{n} a_k = \\ a_i^{n-2} \sum_{j\neq i}^{n} \prod_{k\neq j} a_k + a_i^n \sum_{j\neq i}^{n} \prod_{k\neq j} a_k &= a_i^{n-2}(1 + a_i^2) \sum_{j\neq i}^{n} \prod_{k\neq j} a_k. \end{aligned}$$

To prove iii) we observe first that $n \geq 1$, $k \geq 0, \forall i = 1, \ldots n$:

$$\sum_{j=0}^{n} (-1)^j a_i^{n+k-j} S_j^{(n)}(\mathbf{a}_n) = 0, \tag{1.12}$$

which is elementary to notice. Secondly, using (1.5) we have:

$$\begin{aligned} \sum_{k=0}^{n} (-1)^k S_k^{(n)} B_{n,,m-k} &= A_n \sum_{k=0}^{n} (-1)^k S_k^{(n)} \sum_{i=1}^{n} \frac{a_i^{n+m-k-1}}{\prod_{j\neq i}^{n} (a_i - a_j)(1 - a_i a_j)} \\ &= A_n \sum_{i=1}^{n} \frac{1}{\prod_{j\neq i}^{n} (a_i - a_j)(1 - a_i a_j)} \sum_{k=0}^{n} (-1)^k S_k^{(n)} a_i^{n+m-k-1} = 0, \end{aligned}$$

by (1.12). □

Theorem 2. *For every $n \geq 0$ the family of polynomials orthogonal with respect to f_{Kn} is of the form:*

$$P_m^{(n)}(x|\mathbf{a}_n) = \sum_{j=0}^{n\wedge(2m+2)} (-1)^j U_{m-j}(x) S_j^{(n)}(\mathbf{a}_n), \tag{1.13}$$

$m\geq 0$, *where* $S_j^{(n)}$ *is given by (1.7).*

Proof. Since order of the polynomial P_m must be equal to m and we have the following identity :

$$U_{-m-2}(x)=-U_m(x),$$

both those facts explain why we must have $n\wedge(2m+2)$ as upper summation limit in (1.13).

The fact that polynomials $P_m^{(n)}$ can be expressed as linear combination of the last at most $n+1$ polynomials U_m follows directly from [11](Proposition 1 iii)) or from [3] (Lemma 3.1). Similar fact was noticed for $n=1,2$ earlier by Maroni in his papers published in the 90's. However the form of coefficients of this linear combination can be most easily found arguing by induction.

Assertion of the theorem is true for $n=0,1$. For the proof see e.g. formula (14.18.4) of [6] with $q=0$ or e.g. [15] (3.1 with $q=0$). We see that then $h_m(x|a,0)$ $=U_m(x)-aU_{m-1}(x)$, $m\geq 1$ and $h_0(x|a,0)=1$. Now recall Theorem 1 of [14]. It states that if we have two measures dA and dB related by the formula $dA=\frac{C}{x+D}dB$ and we know polynomials orthogonal $\{b_m\}$ with respect to dB then there exist a numerical sequence κ_m such that $m-$th polynomial d_m orthogonal with respect to dA is equal to $d_m=b_m+\kappa_m b_{m-1}$. By the way the sequence κ_m is defined by the recursion:$\kappa_m=\beta_{m-1}-\hat{\beta}_{m-1}/\kappa_{m-1}+D$, with $\kappa_1=\beta_0+D-C$ where sequences β_m and $\hat{\beta}_m$ are the coefficients of the 3-term recurrence satisfied by polynomials b_m. More precisely $b_{m+1}(x)=(x-\beta_m)b_m(x)-\hat{\beta}_{m-1}b_{m-1}(x)$. Hence by induction assumption we have

$$P_m^{(n)}(x|\mathbf{a}_n)=\sum_{j=0}^{n\wedge(2m+2)}(-1)^jU_{m-j}(x)S_j^{(n)}(\mathbf{a}_n).$$

Further by the quoted Theorem we know that

$$P_m^{(n+1)}(x|\mathbf{a}_{n+1})=P_m^{(n)}(x|\mathbf{a}_n)+\kappa_m^{(n+1)}P_{m-1}^{(n)}(x|\mathbf{a}_n). \tag{1.14}$$

Notice that from (1.6) and (1.11) it follows that $m\geq 1:$

$$\int_{-1}^{1}P_m^{(n)}(x|\mathbf{a}_n)f_{K(n)}(x|\mathbf{a}_n)dx=0.$$

Now let us notice that constant $\kappa_m^{(n+1)}$ can be found from the condition

$$\int_{-1}^{1}P_m^{(n+1)}(x|\mathbf{a}_{n+1})f_{K(n+1)}(x|\mathbf{a}_{n+1})dx=0$$

for all $m\geq 1$.

Using (1.14) we deduce that

$$P_m^{(n+1)}(x|\mathbf{a}_{n+1})=\sum_{j=0}^{n\wedge(2m+2)}(-1)^jU_{m-j}(x)S_j^{(n)}(\mathbf{a}_n)+\kappa_m^{(n+1)}\sum_{j=0}^{n\wedge 2m}(-1)^jU_{m-j-1}(x)S_j^{(n)}(\mathbf{a}_n)$$

$$=U_m(x)-U_{m-1}(x)(S_1^{(n)}(\mathbf{a}_n)-\kappa_m^{(n+1)})+U_{m-2}(x)(S_2^{(n)}(\mathbf{a}_n)-\kappa_m^{(n+1)}S_1^{(n)}(\mathbf{a}_n))...$$

Now we see that for $\kappa_m^{(n+1)}=-a_{n+1}$ we get $S_1^{(n)}(\mathbf{a}_n)-\kappa_m^{(n+1)}=S_1^{(n+1)}(\mathbf{a}_{n+1})$, $S_2^{(n)}(\mathbf{a}_n)-\kappa_m^{(n+1)}S_1^{(n)}(\mathbf{a}_n)=S_2^{(n+1)}(\mathbf{a}_{n+1})$ and so on. Finally $\kappa_m^{(n+1)}S_n^{(n)}(\mathbf{a}_n)=$

$-S_{n+1}^{(n+1)}(\mathbf{a}_{n+1})$. Hence with such choice of $\kappa_m^{(n+1)}$ we get the following formula

$$P_m^{(n+1)}(x|\mathbf{a}_{n+1}) = \sum_{j=0}^{(n+1)\wedge(2m+2)} (-1)^j U_{m-j}(x) S_j^{(n+1)}(\mathbf{a}_{n+1}),$$

which on its side leads to $\int_{-1}^{1} P_m^{(n+1)}(x|\mathbf{a}_{n+1}) f_{K(n+1)}(x|\mathbf{a}_{n+1})dx = 0$ for all $m \geq 1$. ☐

Remark 4. *Notice that polynomials $P_m^{(n)}(x|\mathbf{a}_n)$ are neither monic nor orthonormal.*

Remark 5. *Notice that for $m \geq n$ we have the following 3 term recurrence formula:*

$$2xP_m(x|\mathbf{a}_n) = P_{m+1}(x|\mathbf{a}_n) + P_{m-1}(x|\mathbf{a}_n),$$

which follows directly similar formula to be satisfied by the Chebyshev polynomials $\{U_n\}$. For $m < n$ the problem is more complex. It follows the fact that we naturally have:$P_{-1}(x|\mathbf{a}_n) = 0$, $P_0(x|\mathbf{a}_n) = 1$. However we also have

$$P_1(x|\mathbf{a}_n) = 2x(1 - S_4(\mathbf{a}_n) - S_1(\mathbf{a}_n) + S_3(\mathbf{a}_n),$$

which results in

$$\begin{aligned} P_1(x|a_1) &= 2x - a_1,\ P_1(x|\mathbf{a}_2) = 2x - (a_1 + a_2), \\ P_1(x|\mathbf{a}_3) &= 2x - S_1(\mathbf{a}_3) + S_3(\mathbf{a}_3),\ P_1(x|\mathbf{a}_4) = 2x(1 - S_4(\mathbf{a}_4)) - S_1(\mathbf{a}_4) + S_3(\mathbf{a}_4), \end{aligned}$$

and so on. Note that for $n = 1, 2, 3, 4$ we get polynomials of order 1 of Askey–Wilson scheme for $q = 0$, i.e. big-Hermite ($n = 1$), Al-Salam–Chihara ($n = 2$), continuous dual Hahn ($n = 3$), and Askey–Wilson ($n = 4$).

Similarly for $m = 2$ we have

$$P_2(x|\mathbf{a}_n) = \sum_{j=0}^{6} (-1)^j U_{2-j}(x) S_j(\mathbf{a}_n),$$

which gives

$$\begin{aligned} P_2(x|a_1) &= U_2(x) - a_1 U_1(x),\ P_2(x|\mathbf{a}_2) = U_2(x) - (a_1 + a_2)U_1(x) + a_1 a_2, \\ P_2(x|\mathbf{a}_3) &= U_2(x) - S_1(\mathbf{a}_3)U_1(x) + S_2(\mathbf{a}_3),\ P_2(x|\mathbf{a}_4) = U_2(x) - S_1(\mathbf{a}_4)U_1(x) + S_2(\mathbf{a}_4) - S_4(a_4), \end{aligned}$$

Similarly we notice connection with big-Hermite ($n = 1$), Al-Salam–Chihara ($n = 2$), continuous dual Hahn ($n = 3$), and Askey–Wilson ($n = 4$) polynomials for $q = 0$. Compare [17]*(2.3 with set $q = 0$ and successively $a = d$, $d = 0\&c = 0, d = 0\&c = 0\&b = 0$),*

Proposition 2. *Let $X \sim f_{Kn}(x|\mathbf{a}_n)$ then $\forall\,|t| < 1$:*

$$E\frac{1}{1 + t^2 - 2tX} = \frac{Q_n(t|\mathbf{a}_n)}{\prod_{i=1}^{n}(1 - ta_i)},$$

where

$$Q_n(t|\mathbf{a}_n) = A_n \sum_{i=1}^{n} a_i^{n-1} \prod_{j\neq i}^{n} \frac{(1 - a_j t)}{(a_i - a_j)(1 - a_i a_j)}$$

is a polynomial of degree $\max(n-2, 0)$ *in t.*

Proof. Recall that $\sum_{j=0}^{\infty} t^n U_n(x) = 1/(1+t^2-2tx)$. Thus

$$\int_{-1}^{1} (\sum_{j=0}^{\infty} t^j U_n(x)) f_{Kn}(x|\mathbf{a}_n)dx = A_n \sum_{j=0}^{\infty} t^j \sum_{i=1}^{n} \frac{a_i^{n+j-1}}{\prod_{j\neq i}^{n}(a_i-a_j)(1-a_ia_j)}$$

$$= A_n \sum_{i=1}^{n} \frac{a_i^{n-1}}{\prod_{j\neq i}^{n}(a_i-a_j)(1-a_ia_j)} \sum_{j\geq 0} t^j a_i^j = A_n \sum_{i=1}^{n} \frac{1}{1-a_it} \frac{a_i^{n-1}}{\prod_{j\neq i}^{n}(a_i-a_j)(1-a_ia_j)}$$

$$= \frac{1}{\prod_{i=1}^{n}(1-ta_i)} A_n \sum_{i=1}^{n} a_i^{n-1} \prod_{j\neq i}^{n} \frac{(1-a_jt)}{(a_i-a_j)(1-a_ia_j)}.$$

The fact that Q_n is a polynomial of degree $n-2$ follows the fact that $a_i^{n-1}\prod_{j\neq i}^{n}(1-a_jt)$ is a polynomial of degree $n-1$, but the coefficient by t^{n-1} is equal to $S_n(\mathbf{a}_n)a_i^{n-2}$. The assertion follows from (1.10). □

Remark 6. *We have by direct calculations:*

$$\begin{aligned} Q_1(t|a) &= 1, Q_2(t|\mathbf{a}_2) = 1, Q_3(t|\mathbf{a}_3) = 1 - tS_3(\mathbf{a}_3), \\ Q_4(t|\mathbf{a}_4) &= ((1-S_4) - t(S_3 - S_1S_4) + t^2S_4(1-S_4))/(1-S_4). \end{aligned}$$

As an immediate consequence of this formula (1.3) and definition of $B_{n,k}$ we get characteristic functions of numbers $B_{n,k}(\mathbf{a}_n)$:

Corollary 3. $\forall n \geq 0:$

$$\sum_{k\geq 0} t^k B_{n,k}(\mathbf{a}_n) = Q_n(t|\mathbf{a}_n)/\prod_{i=1}^{n}(1-ta_i).$$

2. Complex parameters

In this section we will study properties of the generalized Kesten-MacKay distributions for even $n = 2k$ and parameters a_i , $i = 1, \ldots, 2k$ being complex and forming conjugate pairs. The new parameters will have new names namely for the conjugate the pair say $a_i = \rho_i \exp(i\theta_i)$ and $a_{k+i-1} = \rho_i \exp(-i\theta_i)$ we will denote $y_i = \cos\theta_i$ for $i = 1, \ldots, k$. Besides notice that we have:

$$\begin{aligned} & (1+\rho_i^2\exp(2i\theta_i) - 2x\rho_i\exp(i\theta_i))(1+\rho_i^2\exp(-2i\theta_i) - 2x\rho_i\exp(-i\theta_i)) \\ = \; & 1+\rho_i^4 + 4x^2\rho_i^2 - 4x\rho_i(1+\rho_i^2)\cos\theta_i + 2\rho_i^2\cos 2\theta_i = w(x, y_i|\rho_i), \end{aligned}$$

where we denoted for simplicity:

$$w(x,y|\rho) = (1-\rho^2)^2 - 4xy\rho(1+\rho^2) + 4\rho^2(x^2+y^2). \tag{2.1}$$

Hence now the density $f_{K2k}(x|a_1, \ldots, a_{2k})$ will have the following form that we will denote by $f_{Mk}(x|\mathbf{y}_k, \boldsymbol{\rho}_k)$:

$$f_{Mk}(x|\mathbf{y}_k, \boldsymbol{\rho}_k) = A_{2k}\frac{2\sqrt{1-x^2}}{\pi\prod_{j=1}^{k} w(x, y_i|\rho_i)}, \tag{2.2}$$

with $|\rho_i| < 1$, $|y_i| \leq 1$.

Following Remark 2 we have:

Lemma 1. *i)* $A_2 = 1-\rho^2$,

ii) $S_1^{(4)}(\mathbf{a}_4) = y_1\rho_1 + y_2\rho_2$, $S_2^{(4)}(\mathbf{a}_4) = \rho_1^2 + \rho_2^2 + 4y_1y_2\rho_1\rho_2$, $S_3^{(4)}(\mathbf{a}_4) = 2\rho_1\rho_2(\rho_1y_1 + \rho_2y_2)$, $A_4 = (1-\rho_1^2)(1-\rho_2^2)w(y_1, y_2|\rho_1\rho_2)/(1-\rho_1^2\rho_2^2)$

iii) $S_1^{(6)}(\mathbf{a}_6) = \rho_1 y_1 + \rho_2 y_2 + \rho_3 y_3$, $S_2^{(6)}(\mathbf{a}_6) = \rho_1^2 + \rho_2^2 + \rho_3^2 + 4(\rho_1\rho_2 y_1 y_2 + \rho_1\rho_3 y_1 y_3$
$+ \rho_2\rho_3 y_2 y_3)$,
$S_3^{(6)}(\mathbf{a}_6) = 2(\rho_1^2 + \rho_3^2)\rho_2 y_2 + 2(\rho_2^2 + \rho_3^2)\rho_1 y_1 + 2(\rho_1^2 + \rho_2^2)\rho_3 y_3 + +8\rho_1\rho_2\rho_3 y_1 y_2 y_3$,
$S_4^{(6)}(\mathbf{a}_6) = \rho_1^2\rho_2^2 + \rho_2^2\rho_3^2 + \rho_1^2\rho_3^2 + 4\rho_1\rho_2\rho_3(\rho_3 y_1 y_2 + \rho_2 y_1 y_3 + \rho_1 y_2 y_3)$,
$S_5^{(6)}(\mathbf{a}_6) = 2\rho_1\rho_2\rho_3(\rho_1\rho_2 y_3 + \rho_1\rho_3 y_2 + \rho_2\rho_3 y_1)$, $S_6^{(6)}(\mathbf{a}_6) = \rho_1^2\rho_2^2\rho_3^2$,
$A_6 = (1-\rho_1^2)(1-\rho_2^2)(1-\rho_3^2)w(y_1, y_2|\rho_1\rho_2)w(y_2, y_3|\rho_2\rho_3)w(y_1, y_3|\rho_1\rho_3)/w3(y_1, y_2, y_3, \rho_1, \rho_2, \rho_3)$,
where we denoted

$$w3(y_1, y_2, y_3, \rho_1, \rho_2, \rho_3) = (1 - \rho_1^2\rho_2^2)(1 - \rho_2^2\rho_3^3)(1 - \rho_1^2\rho_3^2)(1 - \rho_1^2\rho_2^2\rho_3^2) \tag{2.3}$$
$$-4\rho_1\rho_2\rho_3(1 + \rho_1^2\rho_2^2\rho_3^2)(\rho_1(1 - \rho_2^2)(1 - \rho_3^2)y_2 y_3 + \rho_2(1 - \rho_1^2)(1 - \rho_3^2)y_1 y_3 + \rho_3(1 - \rho_1^2)(1 - \rho_2^2)y_1 y_2)$$
$$+4\rho_1^2\rho_2^2\rho_3^2((1 - \rho_1^2)(1 - \rho_2^2\rho_3^2)y_1^2 + (1 - \rho_2^2)(1 - \rho_1^2\rho_3^2)y_2^2 + (1 - \rho_3^2)(1 - \rho_1^2\rho_2^2)y_3^2).$$

Proof. All calculations were done using Mathematica 10. □

Remark 7. *Recall that for all* $|x|, |y| \leq 1$ *and* $|\rho| < 1$ *we have useful expansion:*

$$\frac{1 - \rho^2}{w(x, y|\rho)} = \sum_{j=0}^{\infty} \rho^j U_j(x)U_j(y), \tag{2.4}$$

which is nothing else but the famous Poisson–Mehler formula for $q = 0$ *(for the reference see e.g.* [4] *(13.1.24) or for alternative proof* [11]*)*

Following the above mentioned remark we have:

$$f_{Mk}(x|\mathbf{y}_k, \boldsymbol{\rho}_k) = \frac{2A_{2k}}{\pi\prod_{j=1}^{k}(1 - \rho_j^2)}\sqrt{1 - x^2}\sum_{m_1, m_2, \ldots m_k = 0}^{\infty}\prod_{j=1}^{k}\rho_j^{m_j}U_{m_j}(x)U_{m_j}(y_j), \tag{2.5}$$

$$\prod_{j=1}^{k}(1 - \rho_j^2)/A_{2k} = \sum_{m_1, m_2, \ldots m_k = 0}^{\infty} V_{m_1, \ldots, m_k}\prod_{j=1}^{k}\rho_j^{m_j}U_{m_j}(y_j), \tag{2.6}$$

where as before, above $V_{k_1, \ldots, k_n} = \int_{-1}^{1}\frac{2}{\pi}\sqrt{1 - x^2}\prod_{i=1}^{n}U_{k_i}(x)dx$.

Since each density f_{Mk} can be presented as linear combination of $f_{M1}(x|y_i, \rho_i)$, $k = 1, \ldots, k$ (by simple fraction decomposition) we will analyze f_{M1} first. We have the following result:

Theorem 3. *i)* $\forall y \in [-1, 1] : \int_{-1}^{1} f_{M1}(x|y, \rho) = 1$,
ii) $\int_{-1}^{1} f_{M1}(x|y, \rho)\frac{2}{\pi}\sqrt{1 - y^2}dy = \frac{2}{\pi}\sqrt{1 - x^2}$,
iii) $\int_{-1}^{1} f_{M1}(x|y_1, \rho_1)f_{M1}(y_1|y_2, \rho_2)dy_1 = f_{M1}(x|y_2, \rho_1\rho_2)$.

Proof. i) Either we use directly properties of $f_{K2}(x|a, b)$ and the fact that in our case $ab = \rho^2$, or we apply (2.4). ii) follows directly from i) and the fact that $w(x, y|\rho) = w(y, x|\rho)$. iii) we have:

$$\int_{-1}^{1} f_{M1}(x|y_1, \rho_1)f_{M1}(y_1|y_2, \rho_2)dy_1 =$$
$$\frac{2}{\pi}\sqrt{1 - x^2}\int_{-1}^{1}\frac{2}{\pi}\frac{(1 - \rho_1^2)(1 - \rho_2^2)\sqrt{1 - y_1^2}}{w(y_1, x|\rho_1)w(y_1, y_2|\rho_2)}dy_1 =$$
$$\frac{2}{\pi}(1 - \rho_1^2)(1 - \rho_2^2)\sqrt{1 - x^2}/A_4 = f_{M2}(x|y_2, \rho_1\rho_2),$$

since $A_4 = (1 - \rho_1^2)(1 - \rho_2^2)w(x, y_2|\rho_1\rho_2)/(1 - \rho_1^2\rho_2^2)$. □

Remark 8. *Results of the above Theorem indicate possible applications of the distributions f_{M1} and Wigner in multivariate analysis and stochastic processes. More precisely assertion i) shows that $f_{M1}(x|y,\rho)$ is in fact a conditional distribution. ii) shows that $f_{M1}(x|y,\rho)f_{M0}(y)$ can be treated as a density of certain bivariate distribution with f_{M0} that is Wigner distribution as its marginals. finally iii) is nothing else but the so-called Chapman–Kolmogorov property. These properties are known and applied in stochastic processes, see e.g.* [2] *and* [12]. *We quoted them for the sake of completeness of the paper and also in order to present new proofs of these properties directly basing on the general properties of generalized Kesten distributions discussed in the first part of this paper.*

Following the above mentioned remark let us denote by $f_2(x,y|\rho)$ denote $2-$dimensional measure defined by:

$$f_{2M}(x,y|\rho)=f_{M1}(x|y,\rho)f_{M0}(y)=\frac{(1-\rho^2)\sqrt{(1-x^2)(1-y^2)}}{\pi^2 w(x,y|\rho)}. \tag{2.7}$$

Remark 9. *Notice also that*

$$\begin{aligned} f_{M2}(x|y_1,\rho_1,y_2,\rho_2) &= \frac{2}{\pi}\frac{\sqrt{1-x^2}(1-\rho_1^2)(1-\rho_2^2)w(y_1,y_2|\rho_1\rho_2)}{w(x,y_1|\rho_1)w(x,y_2|\rho_2)(1-\rho_1^2\rho_2^2)} \\ &= \frac{f_{M1}(y_1|x,\rho_1)f_{M1}(x|y_2,\rho_2)f_{M0}(y_2)}{f_{M1}(y_1|y_2,\rho_1\rho_2)f_{M0}(y_2)}, \end{aligned}$$

which can be interpreted in the following way. Let us consider 3 element discrete Markov chain X_1,X_2,X_3 such that transition density $X_2|X_3$ is $f_{M1}(x|y_2,\rho_2)$, transition $X_1|X_2$ is $f_{M1}(x|y_1,\rho_1)$ while marginal density of X_3 is $f_{M0}(y_2)$ then the conditional density of $X_2|X_1,X_3$ is $f_{M2}(x|y_1,\rho_1,y_2,\rho_2)$.

Lemma 2. $\int_{-1}^{1}\frac{2}{\pi}\sqrt{1-y_1^2}\frac{w3(y_1,y_2,y_3,\rho_1,\rho_2,\rho_3)}{w(y_1,y_2|\rho_1\rho_2)w(y_2,y_3|\rho_2\rho_3)w(y_1,y_3|\rho_1\rho_3)}dy_1=\frac{1-\rho_2^2\rho_3^2}{w(y_2,y_3,|\rho_2\rho_3)}$.

Proof. We start from (2.5) considered for $k=3$ and get $\prod_{j=1}^{3}(1-\rho_i^2)/A_6=\frac{w3(y_1,y_2,y_3,\rho_1,\rho_2,\rho_3)}{w(y_1,y_2|\rho_1\rho_2)w(y_2,y_3|\rho_2\rho_3)w(y_1,y_3|\rho_1\rho_3)}=$ $\sum_{m_1=0}^{\infty}\sum_{m_2=0}^{\infty}\sum_{m_3=0}^{\infty}\rho_1^{m_1}\rho_2^{m_2}\rho_3^{m_3}V_{m_1m_2m_3}U_{m_1}(y_1)U_{m_2}(y_2)U_{m_3}(y_3)$ basing on Lemma 1. Now $\int_{-}^{1}\frac{2}{\pi}\sqrt{1-y_1^2}\frac{w3(y_1,y_2,y_3,\rho_1,\rho_2,\rho_3)}{w(y_1,y_2|\rho_1\rho_2)w(y_2,y_3|\rho_2\rho_3)w(y_1,y_3|\rho_1\rho_3)}dy_1=$ $\sum_{m_2=0}^{\infty}\sum_{m_3=0}^{\infty}\rho_2^{m_2}\rho_3^{m_3}V_{0m_2m_3}U_{m_2}(y_2)U_{m_3}(y_3)=\sum_{m_2=0}^{\infty}(\rho_2\rho_3)^{m_2}U_{m_2}(y_2)U_{m_2}(y_3)$ $=\frac{1-\rho_2^2\rho_3^2}{w(y_2,y_3,|\rho_2\rho_3)}$. □

From this Lemma we derive the following important conclusion. Namely that the function:

$$g(y_1,y_2,y_3|\rho_1,\rho_2,\rho_3)= \tag{2.8}$$
$$\frac{8}{\pi^3}\sqrt{1-y_1^2}\sqrt{1-y_2^2}\sqrt{1-y_3^2}\frac{w3(y_1,y_2,y_3,\rho_1,\rho_2,\rho_3)}{w(y_1,y_2|\rho_1\rho_2)w(y_2,y_3|\rho_2\rho_3)w(y_1,y_3|\rho_1\rho_3)}$$

can be treated as the density of some $3D$ distribution with $2D$ marginals equal to $f_{2M}(y_1,y_2|\rho_1\rho_2)$, $f_{2M}(y_1,y_3|\rho_1\rho_3)$, $f_{2M}(y_2,y_3|\rho_2\rho_3)$.

REFERENCES

[1] Bożejko, Marek; Kümmerer, Burkhard; Speicher, Roland. q-Gaussian processes: non-commutative and classical aspects. *Comm. Math. Phys.* **185** (1997), no. 1, 129–154. MR1463036 (98h:81053)

[2] Bryc, Włodzimierz; Matysiak, Wojciech; Szabłowski, Paweł J. Probabilistic aspects of Al-Salam-Chihara polynomials. *Proc. Amer. Math. Soc.* **133** (2005), no. 4, 1127–1134 (electronic). MR2117214 (2005m:33033)

[3] Delgado, Antonia M.; Geronimo, Jeffrey S.; Iliev, Plamen; Xu, Yuan. On a two-variable class of Bernstein-Szegő measures. *Constr. Approx.* **30** (2009), no. 1, 71–91. MR2519655

[4] Ismail, Mourad E. H. Classical and quantum orthogonal polynomials in one variable. With two chapters by Walter Van Assche. With a foreword by Richard A. Askey. Encyclopedia of Mathematics and its Applications, 98. *Cambridge University Press,* Cambridge, 2005. xviii+706 pp. ISBN: 978-0-521-78201-2; 0-521-78201-5 MR2191786 (2007f:33001)

[5] Kesten, Harry. Symmetric random walks on groups. *Trans. Amer. Math. Soc.* **92** 1959 336–354. MR0109367 (22 #253)

[6] Koekoek, Roelof; Lesky, Peter A.; Swarttouw, René F. Hypergeometric orthogonal polynomials and their q-analogues. With a foreword by Tom H. Koornwinder. *Springer Monographs in Mathematics. Springer-Verlag*, Berlin, 2010. xx+578 pp. ISBN: 978-3-642-05013-8 MR2656096 (2011e:33029)

[7] Oren, Idan; Godel, Amit; Smilansky, Uzy. Trace formulae and spectral statistics for discrete Laplacians on regular graphs. I. *J. Phys. A* **42** (2009), no. 41, 415101, 20 pp. MR2545624 (2010k:05181)

[8] Oren, Idan; Smilansky, Uzy. Trace formulas and spectral statistics for discrete Laplacians on regular graphs (II). *J. Phys. A* **43** (2010), no. 22, 225205, 13 pp. MR2647344 (2011g:81085)

[9] McKay, Brendan D. The expected eigenvalue distribution of a large regular graph. *Linear Algebra Appl.* **40** (1981), 203–216. MR0629617 (84h:05089)

[10] Sodin, Sasha. Random matrices, nonbacktracking walks, and orthogonal polynomials. *J. Math. Phys.* **48** (2007), no. 12, 123503, 21 pp. MR2377835 (2008m:60082)

[11] Szabłowski, Paweł J. Expansions of one density via polynomials orthogonal with respect to the other. *J. Math. Anal. Appl.* **383** (2011), no. 1, 35–54. MR2812716, http://arxiv.org/abs/1011.1492

[12] Szabłowski, Paweł J. $q-$Wiener and $(\alpha,q)-$ Ornstein–Uhlenbeck processes. A generalization of known processes, *Theory of Probability and Its Applications,* **56** (4), 2011, 742–772, http://arxiv.org/abs/math/0507303

[13] Paweł J Szabłowski, Askey-Wilson integral and its generalizations, *Advances in Difference Equations* 2014, **2014:316**

[14] Szabłowski, Paweł J. On affinity relating two positive measures and the connection coefficients between polynomials orthogonalized by these measures. *Appl. Math. Comput.* **219** (2013), no. 12, 6768–6776. MR3027843

[15] Szabłowski, Paweł, J. On the $q-$Hermite polynomials and their relationship with some other families of orthogonal polynomials, *Dem. Math.* **66(**2013) no.4, 679-708, http://arxiv.org/abs/1101.2875,

[16] Szabłowski, Paweł, J., Moments of $q-$Normal and conditional $q-$Normal distributions, *Stoch. and Porob. Letters,* **106** (2015), 65-72

[17] Szabłowski, Paweł J. Befriending Askey–Wilson polynomials, *Infin. Dimens. Anal. Quantum Probab. Relat. Top. ,* Vol . **17**, No. 3 (2014) 1450015 (25 pages), http://arxiv.org/abs/1111.0601.

Warsaw University of Technology,, Department of Mathematics and Information Sciences

E-mail address: `pawel.szablowski@gmail.com`